

JOSEPHI NICOLAI
LAURENTI
AUSTRIACI VIENNENSIS
SPECIMEN
MEDICUM,
EXHIBENS
SYNOPSISIN REPTILIIUM
EMENDATAM
CUM EXPERIMENTIS CIRCA VENENA
ET ANTIDOTA REPTILIIUM AUSTRIACORUM.



VIENNÆ,
TYP. JOAN. THOM. NOB. DE TRATTNERN,
CÆS. REG. AULÆ TYPOGR. ET BIBLIOP.

1768.

CIV. *Seps sericeus*. *Tab. II. fig. 5.*

DIAGN.

Collario adnato; scutis intermediis quadratis; cute laxa, delicatissima, tenerrima, subsquammulata, fusca.

Habitat in stratis lapideis circa aquas.

CV. *Seps argus*. *Tab. I. fig. 5.*

DIAGN.

Collario adnato: scutis intermediis quadratis; corpore quaquaversum ocellato.

Habitat in muris verticalibus.

CVI. *Seps muralis*. *Tab. I. fig. 4.*

DIAGN.

Collario adnato; scutis intermediis quadratis; femoribus transversis; corpore humili; tænia utrinque ad latera thoracis saturatiori.

Habitat in mœnibus, ruderibus, aggeribus lapidosis passim ubique.

β. Collarii squammæ infimæ solutæ; scuta seriei intermediæ subrotunda.

CVII. *Seps terrestris*. *Tab. III. fig. 1.*

DIAGN.

Collario soluto; scutis intermediis subrotundis; corpore fusco; utrinque serie macularum obsoletarum.

Habitat in terra lapidosa plana.